

ParaGard[®] Allwormer Tablets for Small Dogs and Puppies

4
TABLETS

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

ParaGard[®] Allwormer Tablets for Small Dogs and Puppies

EACH TABLET CONTAINS 25 mg PRAZIQUANTEL, 112.5 mg OXIBENDAZOLE

For the treatment and control of Roundworm, Hookworm,
Whipworm and Tapeworm, including hydatids, in dogs.
Does not control heartworm in dogs. Palatable liver flavour.



C1835

ParaGard[®] Allwormer Tablets for Small Dogs and Puppies

4
TABLETS

1P6331-2



SAFETY DIRECTIONS
May irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID
If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

DISPOSAL
Dispose of empty container by wrapping with paper and putting in garbage.

STORAGE
Store below 30°C (Room Temperature).

Boehringer Ingelheim Animal Health Australia Pty Ltd
Level 1, 78 Waterloo Road, North Hyde NSW 2113
solutions on 1800 808 691.
APVMA Approval No.: 80477/100937
@ ParaGard is a registered trademark of the Boehringer Ingelheim Group.



4
TABLETS

ParaGard[®] Allwormer Tablets
for Small Dogs and Puppies

Indications
For use against the following species of worms: *Toxascaris leonina*, *Toxocara canis*, *Ancylostoma caninum*, *Uncinaria stenocephala*, *Trichostrongylus axei*, *Dipylidium caninum*, *Taenia ovis*, *Taenia pisiformis*, *Taenia hydatigena* and *Echinococcus granulosus* (hydatids) in dogs.

If worm problems persist, consult a veterinarian.

DIRECTIONS FOR USE
Puppies: Treat according to weight at 2, 4, 8 and 12 weeks - thereafter every 3 months.
Adult Dogs: Treat every 3 months.
Breeding Bitch: Treat during the mating period, during the gestation period one week before birth and 3 to 4 weeks after birth.
Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks. Where whipworm is a problem, treat every 6-8 weeks after 3 months of age.

Dosage and Administration
1 tablet/5 kg body weight

PARAGARD ALLWORMER is to be used as a single dose directly as a treat or crushed and put on food.

Fleas are intermediate hosts for tapeworm (*Dipylidium caninum*). Effective flea control will reduce the incidence of tapeworm.

ParaGard[®] Allwormer Tablets for Medium Dogs

4
TABLETS

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

ParaGard[®] Allwormer Tablets for Medium Dogs

EACH TABLET CONTAINS 50 mg PRAZIQUANTEL, 225 mg OXIBENDAZOLE

For the treatment and control of **Roundworm, Hookworm, Whipworm and Tapeworm**, including **hydatids**, in dogs.
Does not control heartworm in dogs. Palatable **liver flavour**.



C1836

ParaGard[®] Allwormer Tablets for Medium Dogs

4
TABLETS

1P6333-2



@ Paragard is a registered trademark of the Boehringer Ingelheim Group.
APVMA Approval No.: 80478/100938

For more information, contact Boehringer Ingelheim customer solutions on 1800 808 691.
Boehringer Ingelheim Animal Health Australia Pty Ltd
Level 1, 78 Waterloo Road, North Ryde NSW 2113
Store below 30°C (Room Temperature).

STORAGE

putting in garbage.
Dispose of empty container by wrapping with paper and

DISPOSAL

Information Centre, Phone Australia 131126.

FIRST AID

If poisoning occurs contact a doctor or Poisons

SAFETY DIRECTIONS

May irritate the eyes. Avoid contact with eyes. Wash hands after use.



Indications
For use against the following species of worms: *Toxascaris leonina*, *Toxocara canis*, *Ancylostoma caninum*, *Uncinaria stenocephala*, *Trichuris vulpis*, *Dipylidium caninum*, *Uncinaria stenocephala*, *Trichuris vulpis*, *Dipylidium caninum*, *Uncinaria stenocephala*, *Taenia hydatigena* and *Echinococcus granulosus* (hydatids) in dogs.
If worm problems persist, consult a veterinarian.

Directions for Use
Puppies: Treat according to weight at 2, 4, 8 and 12 weeks after 3 months of age.
Adult Dogs: Treat every 3 months.
Breeding Bitch: Treat during the mating period, during the gestation period one week before birth and 3 to 4 weeks after birth.
Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks. Where whipworm is a problem, treat every 6-8 weeks after 3 months of age.

PARAGARD ALLWORMER is to be used as a single dose treatment. Tablets are palatable and may be administered directly as a treat or crushed and put on food.
Effective flea control will reduce the incidence of tapeworm. Fleas are intermediate hosts for tapeworm (*Dipylidium caninum*).

4
TABLETS

ParaGard[®] Allwormer Tablets
for Medium Dogs

ParaGard® Allwormer Tablets for Large Dogs

3
TABLETS

CAUTION
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING
FOR ANIMAL TREATMENT ONLY

ParaGard® Allwormer Tablets for Large Dogs

EACH TABLET CONTAINS 100 mg PRAZQUANTEL, 450 mg OXIBENDAZOLE

For the treatment and control of **Roundworm, Hookworm, Whipworm and Tapeworm**, including **hydatids**, in dogs.
Does not control heartworm in dogs. Palatable liver flavour.



C1837

ParaGard® Allwormer Tablets for Large Dogs

3
TABLETS

1P6335-2



For more information, contact Boehringer Ingelheim customer solutions on 1800 808 691.
Boehringer Ingelheim Animal Health Australia Pty Ltd
Level 1, 78 Waterloo Road, North Hyde NSW 2113
Store below 30°C (Room Temperature).
STORAGE
Dispose of empty container by wrapping with paper and putting in garbage.



SAFETY DIRECTIONS
May irritate the eyes. Avoid contact with eyes. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131126.

DISPOSAL

Dispose of empty container by wrapping with paper and putting in garbage.

Indications
For use against the following species of worms: *Toxascaris leonina*, *Toxocara canis*, *Angiostrongylus cantonum*, *Uncinaria stenocephala*, *Trichostrongylus axei*, *Dipylidium caninum*, *Taenia ovis*, *Taenia pisiformis*, *Taenia hydatigena* and *Echinococcus granulosus* (hydatids) in dogs.
If worm problems persist, consult a veterinarian.

DIRECTIONS FOR USE
Dosage and Administration
1 tablet/20 kg body weight
Puppies: Treat according to weight at 2, 4, 8 and 12 weeks - thereafter every 3 months.
Adult Dogs: Treat every 3 months.
Breeding Bitch: Treat during the mating period, during the gestation period one week before birth and 3 to 4 weeks after birth.
Dogs should not be fed, or allowed to feed on, offal from any species. Dogs in hydatid areas should be treated every 6 weeks. Where whipworm is a problem, treat every 6-8 weeks after 3 months of age.
PARAGARD ALLWORMER is to be used as a single dose treatment. Tablets are palatable and may be administered directly as a treat or crushed and put on food.
Effective flea control will reduce the incidence of tapeworm. Fleas are intermediate hosts for tapeworm (*Dipylidium caninum*).

3
TABLETS

ParaGard® Allwormer Tablets
for Large Dogs