

CAUTION

KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS
FOR ANIMAL TREATMENT ONLY

NexGard SPECTRA® Chewables for Dogs

Each chewable contains 9.4 mg Afoxolaner and 1.9 mg Milbemycin Oxime (2 – 3.5 kg Dogs).

Each chewable contains 18.8 mg Afoxolaner and 3.8 mg Milbemycin Oxime (3.6 – 7.5 kg Dogs).

Each chewable contains 37.5 mg Afoxolaner and 7.5 mg Milbemycin Oxime (7.6 – 15 kg Dogs).

Each chewable contains 75 mg Afoxolaner and 15 mg Milbemycin Oxime (15.1 – 30 kg Dogs).

Each chewable contains 150 mg Afoxolaner and 30 mg Milbemycin Oxime (30.1 – 60 kg Dogs).

- Rapidly kills fleas
- Treats and controls ticks (Paralysis ticks, Brown Dog ticks, Bush ticks)
- Prevents heartworm disease
- Treats and controls roundworms, hookworms and whipworms
- Controls flea allergy dermatitis
- Monthly treatment
- Tasty chew

Indications

For the treatment and prevention of flea infestations, control of ticks, prevention of heartworm and treatment and control of intestinal worms in dogs and puppies from 8 weeks of age.

NexGard Spectra rapidly kills adult fleas and is indicated for the treatment and prevention of flea (*Ctenocephalides* spp.) infestations, for the control of newly acquired adult fleas and for the control of flea allergy dermatitis. Following administration, fleas are killed within 6-8 hours. Rapid onset of action kills newly acquired fleas before they can lay eggs. NexGard Spectra treats and controls Paralysis ticks (*Ixodes holocyclus*), Brown Dog ticks (*Rhipicephalus sanguineus*) and Bush ticks (*Haemaphysalis longicornis*) for one month. NexGard Spectra provides effective control of pre-existing Paralysis tick infestations within 24 hours. *Ixodes holocyclus* ticks do not occur in Western Australia. In tick season, daily searching for, and removal of, any ticks found is recommended.

NexGard Spectra prevents heartworm disease caused by *Dirofilaria immitis* and treats and controls roundworms (*Toxocara canis* and *Toxascaris leonina*), hookworms (*Ancylostoma caninum*) and whipworms (*Trichuris vulpis*). Consult a veterinarian if worm problems persist. Monthly administration is recommended.

DIRECTIONS FOR USE

Precautions

The safe use of NexGard Spectra Chewables in breeding males and females has not been evaluated.

Dosage and Administration

- The recommended dosage for 2 – 3.5 kg dogs is one 0.5 g chewable (9.4 mg afoxolaner and 1.9 mg milbemycin oxime/chewable) to deliver a minimum of 2.5 mg of afoxolaner per kg bodyweight and 0.5 mg of milbemycin oxime per kg of bodyweight.
- The recommended dosage for 3.6 – 7.5 kg dogs is one 1 g chewable (18.8 mg afoxolaner and 3.8 mg milbemycin oxime/chewable) to deliver a minimum of 2.5 mg of afoxolaner per kg bodyweight and 0.5 mg of milbemycin oxime per kg of bodyweight.

- The recommended dosage for 7.6 – 15 kg dogs is one 2 g chewable (37.5 mg afoxolaner and 7.5 mg milbemycin oxime/chewable) to deliver a minimum of 2.5 mg of afoxolaner per kg bodyweight and 0.5 mg of milbemycin oxime per kg of bodyweight.
- The recommended dosage for 15.1 – 30 kg dogs is one 4 g chewable (75 mg afoxolaner and 15 mg milbemycin oxime/chewable) to deliver a minimum of 2.5 mg of afoxolaner per kg bodyweight and 0.5 mg of milbemycin oxime per kg of bodyweight.
- The recommended dosage for 30.1 – 60 kg dogs is one 8 g chewable (150 mg afoxolaner and 30 mg milbemycin oxime/chewable) to deliver a minimum of 2.5 mg of afoxolaner per kg bodyweight and 0.5 mg of milbemycin oxime per kg of bodyweight.

Offer the chewable(s) appropriate for your dog's weight at monthly intervals. Remove only one chewable at a time from the foil-backed blister card. NexGard Spectra may be broken into pieces and fed to dogs who normally swallow treats whole. NexGard Spectra is a soft, highly palatable chewable and is readily consumed by dogs when offered as a treat. Alternatively, NexGard Spectra may be added intact to a small amount of dog food.

Care should be taken that the dog consumes the complete dose, and treated dogs should be observed for a few minutes to ensure that part of the dose is not lost or rejected. If it is suspected that any of the dose has been lost, treat again with another full dose. If a dose is missed, administer the NexGard Spectra chewable(s) appropriate for your dog's weight and resume a monthly dosing regimen. The recommended minimum monthly dosage is 2.5 mg afoxolaner per kg bodyweight and 0.5 mg milbemycin oxime per kg bodyweight:

Dog Size	Dog Weight (kg)	Chewable size (g)	Afoxolaner (mg/chew)	Milbemycin oxime (mg/chew)
Very small	2 – 3.5	0.5	9.4	1.9
Small	3.6 – 7.5	1	18.8	3.8
Medium	7.6 – 15.0	2	37.5	7.5
Large	15.1 – 30.0	4	75.0	15.0
Very Large	30.1 – 60	8	150.0	30.0

For dogs over 60 kg, administer the appropriate combination of chewables. For optimal performance, treat your dog on or about the same day each month.

General Directions

As NexGard Spectra is an oral treatment, there are no restrictions on swimming, bathing and shampooing before or after administration.

Safety to Dogs

NexGard Spectra was shown to be safe in various breeds of dogs (including Collie breeds) and in puppies 8 weeks and older. Afoxolaner is part of the isoxazoline family of chemicals. Adverse reactions to this family of chemicals are rarely observed but may include vomiting, diarrhoea, lethargy, inappetence, itching and very rarely, seizures. Most adverse reactions are self-limiting and of short duration. If you have any concerns, please speak to your veterinarian.

Flea Treatment and Prevention

NexGard Spectra demonstrated rapid control of fleas for a full month. Because fleas are killed rapidly, NexGard Spectra controls flea allergy dermatitis.

Treatment with NexGard Spectra may begin at any time of the year. Dogs become infested with fleas that originate from environmental infestation sites in the home or outdoors. Treatment and prevention of flea infestations is achieved by killing fleas on the dog before they can lay eggs and replenish an environmental infestation. If a severe environmental infestation exists, fleas can continue to emerge into the surroundings for weeks to months, because of the continuing development of flea eggs, larvae and pupae already present at the time of treatment. It may therefore take more than one monthly treatment to achieve control of moderate to heavy pre-existing flea infestations.

In areas where fleas are common year-round, monthly treatment with NexGard Spectra should continue throughout the year without interruption. To minimise the likelihood of flea reinfestation, it is important to treat all animals within a household with an approved flea control product.

Tick Control

For the most effective control of Paralysis ticks (*Ixodes holocyclus*) and to minimise the risk of tick paralysis, NexGard Spectra should be used in conjunction with daily searching and removal of ticks. Treatment with NexGard Spectra alone will not entirely remove the risk of tick paralysis. Speak with your veterinarian for advice on tick searching and removal techniques.

NexGard Spectra demonstrated >95% efficacy for a full month against Paralysis ticks. Brown Dog ticks and Bush ticks. In Australian studies, Paralysis ticks were killed (>95% efficacy) within 48 hours. Brown Dog ticks and Bush ticks were killed (>95% efficacy) within 48 hours.

Treatment with NexGard Spectra may begin at any time of the year. Dogs become infested with ticks that originate from sites which are primarily outdoors. In areas where ticks are common year-round, monthly treatment with NexGard Spectra should continue throughout the year without interruption.

Heartworm Disease Prevention

NexGard Spectra should be administered monthly to prevent heartworm disease.

Treatment with NexGard Spectra should continue throughout the year without interruption. When replacing another heartworm preventative product, the first dose of NexGard Spectra should be administered within one month of the last dose of the previous monthly medication.

Consult your veterinarian before starting a heartworm prevention program. Heartworms can have serious effects on the health of dogs. The use of NexGard Spectra in heartworm positive dogs has not been evaluated.

Gastrointestinal Worms

Roundworms, hookworms and whipworms cause illness in dogs. Regular treatment with NexGard Spectra provides effective treatment and control of roundworms (*T. canis* and *T. leonina*), hookworms (*A. caninum*) and whipworms (*T. vulpis*).

SAFETY DIRECTIONS

Do not swallow. Do not open inner pouch until ready for use. Wash hands after use.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia 131 126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

ADDITIONAL USER SAFETY INFORMATION

Accidental ingestion of afoxolaner can be harmful to children. To avoid accidental ingestion administer the chewable to the dog immediately after removal from the package. Keep out of reach of children. Additional information is in the Safety Data Sheet.

DISPOSAL

Dispose of empty pack by wrapping in paper and placing in garbage.

STORAGE

Store below 30°C (Room Temperature). Protect from light.

MERIAL Australia Pty Ltd
Level 2.01, 16 Giffnock Avenue
North Ryde NSW 2113

For product information and technical enquiries visit www.merial.com.au or call MERIAL Customer Solutions Australia: 1800 808 691.

APVMA Approval Nos:

81008/102149
81010/102155
81011/102157
81012/102161
81055/102246



NexGard Spectra is a registered trademark.
© 2017 MERIAL Limited. P16742